

Clusters of matter and antimatter

I.N. Mishustin¹

*Kurchatov Institute, Russian Research Center, Moscow 123182, Russia
Niels Bohr Institute, University of Copenhagen, DK-2100 Copenhagen, Denmark
Institut für Theoretische Physik, Goethe Universität, D-60054 Frankfurt am Main,
Germany*

Abstract

In this talk I first give a short overview of antinuclei production in recent experiments at RHIC. Then I discuss the possibility of producing new types of nuclear systems by implanting an antibaryon into ordinary nuclei. The structure of nuclei containing one antiproton or antilambda is investigated within the framework of a relativistic mean-field model. Self-consistent calculations predict an enhanced binding and considerable compression in such systems as compared with normal nuclei. I present arguments that the life time of such nuclei with respect to the antibaryon annihilation might be long enough for their observation. Few experimental signatures are suggested. Possible formation of multi-quark-antiquark clusters is also discussed.

1 Introduction

Relativistic heavy-ion collisions open unique possibility of studying antimatter production in the laboratory. Experiments with colliding Au beams at highest energy available at present are conducted at the Relativistic Heavy-Ion Collider (RHIC) in Brookhaven National Laboratory. More than 1000 particles per unit rapidity are produced in a central Au+Au collision at $\sqrt{s} = 130$ AGeV. Most of them are pions, but many baryons and antibaryons are produced too. Moreover, light nuclei and antinuclei have also been registered! As reported by the STAR experiment [1], the rapidity densities of different species at $y \approx 0$ are

$$\frac{dN}{dy} = 32 \text{ (p)}, 24 \text{ (}\bar{\text{p}}\text{)}, 2.7 \cdot 10^{-3} \text{ (}\bar{\text{d}}\text{)}, 6.3 \cdot 10^{-6} \text{ (}^3\bar{\text{He}}\text{)} . \quad (1)$$

¹This work was supported in part by the Deutsche Forschung Gemeinschaft (DFG) under grant 436 RUS 113/711/0-1, and the Russian Fond of Fundamental Research (RFFR) under grant 03-02-04007.

No ${}^4\overline{\text{He}}$ have been observed yet, although their detection is feasible. The corresponding ratios are

$$\frac{\overline{p}}{p} \simeq 0.75, \quad \frac{\overline{d}}{\overline{p}} \simeq 10^{-4}, \quad \frac{{}^3\overline{\text{He}}}{\overline{d}} \simeq 2 \cdot 10^{-3}. \quad (2)$$

This is almost baryon-symmetric state of matter, similar to the one existed in the early Universe around the deconfinement transition.

The BRAHMS experiment has measured rapidity densities of p and $\overline{p}$ in a wider rapidity interval, $0 < y < 3$ [2]. Using these data one finds that the total number of antibaryons produced in a central Au+Au collision is about 200. This is sufficient to make an anti-Pb nucleus! But this is impossible because these antibaryons are widely dispersed in the phase space. Only lightest antinuclei ($\overline{d}$, $\overline{t}$, ${}^3, {}^4\overline{\text{He}}$, ...) can be formed predominantly via the coalescence mechanism.

One may speculate that more exotic clusters of baryons and antibaryons, like $\overline{B} + B$, $\overline{B} + d$, $\overline{B} + \alpha$ or even $\overline{d} + d$, can be formed in ultrarelativistic heavy-ion collisions. If their life times are long enough, they will show up themselves by the delayed annihilation. More heavy clusters like $\overline{B} + A$ where A stands for normal nuclei (${}^4\text{He}$, ${}^{16}\text{O}$, ${}^{40}\text{Ca}$, ...) can be formed by using high-energy antiproton beams. Below I discuss theoretical predictions on properties of such clusters and their possible signatures.

2 Relativistic description of nuclear matter

It has been noticed already many years ago (see e.g. ref. [3]) that nuclear physics may provide a unique laboratory for investigating the Dirac picture of vacuum. The basis for this is given by relativistic mean-field models which are widely used now for describing nuclear matter and finite nuclei. Within this approach nucleons are described by the Dirac equation coupled to scalar and vector meson fields. Scalar S and vector V potentials generated by these fields modify plane-wave solutions of the Dirac equation as follows

$$E^\pm(\mathbf{p}) = V \pm \sqrt{\mathbf{p}^2 + (m - S)^2}. \quad (3)$$

The $+$ sign corresponds to nucleons with positive energy $E_N(\mathbf{p}) = E^+(\mathbf{p}) = V + \sqrt{\mathbf{p}^2 + (m - S)^2}$, and the $-$ sign corresponds to antinucleons with energy $E_{\overline{N}}(\mathbf{p}) = -E^-(-\mathbf{p}) = -V + \sqrt{\mathbf{p}^2 + (m - S)^2}$. It is remarkable that changing sign of the vector potential for antinucleons is exactly what is expected from the G-parity transformation of the nucleon potential. As follows from eq. (3), the spectrum of single-particle states of the Dirac equation in nuclear environment is modified in two ways. First, the mass gap between positive- and negative-energy states, $2(m - S)$, is reduced due to the scalar potential and second, all states are shifted upwards due to the vector potential.

It is well known from nuclear phenomenology that good description of nuclear ground state is achieved with $S \simeq 350$ MeV and $V \simeq 300$ MeV so that the net potential for nucleons with $p \approx 0$ is $V - S \simeq -50$ MeV. Using the same values one obtains for antinucleons a very deep potential, $-V - S \simeq -650$ MeV. Such a potential would produce many strongly bound states in the Dirac sea. However, if these states are occupied they are hidden from the direct observation. Only creating a hole in this sea, i.e. inserting a real antibaryon into the nucleus, would produce an observable effect. If this picture

is correct one should expect the existence of strongly bound states of antinucleons with nuclei [4].

The structure of such systems is calculated using several versions of the relativistic mean-field model (RMF): TM1 [5], NL3 and NL-Z2 [6]. Their parameters were found by fitting binding energies and charge form-factors of spherical nuclei from ^{16}O to ^{208}Pb . The general Lagrangian of the RMF model is written as

$$\begin{aligned}
\mathcal{L} = & \sum_{j=B,\bar{B}} \bar{\psi}_j (i\gamma^\mu \partial_\mu - m_j) \psi_j \\
& + \frac{1}{2} \partial^\mu \sigma \partial_\mu \sigma - \frac{1}{2} m_\sigma^2 \sigma^2 - \frac{b}{3} \sigma^3 - \frac{c}{4} \sigma^4 \\
& - \frac{1}{4} \omega^{\mu\nu} \omega_{\mu\nu} + \frac{1}{2} m_\omega^2 \omega^\mu \omega_\mu + \frac{d}{4} (\omega^\mu \omega_\mu)^2 \\
& - \frac{1}{4} \vec{\rho}^{\mu\nu} \vec{\rho}_{\mu\nu} + \frac{1}{2} m_\rho^2 \vec{\rho}^\mu \vec{\rho}_\mu \\
& + \sum_{j=B,\bar{B}} \bar{\psi}_j (g_{\sigma j} \sigma + g_{\omega j} \omega^\mu \gamma_\mu + g_{\rho j} \vec{\rho}^\mu \gamma_\mu \vec{\tau}_j) \psi_j
\end{aligned} \tag{4}$$

plus Coulomb part. Here summation includes valence baryons B , i. e. the nucleons forming a nucleus, and valence antibaryons $\bar{B}$ inserted in the nucleus. They are treated as Dirac particles coupled to the scalar-isoscalar (σ), vector-isoscalar (ω) and vector-isovector ($\vec{\rho}$) meson fields. The calculations are carried out within the mean-field approximation where the meson fields are replaced by their expectation values. Also a "no-sea" approximation is used. This implies that all occupied states of the Dirac sea are "integrated out" so that they do not appear explicitly. It is assumed that their effect is taken into account by nonlinear terms in the meson Lagrangian. Most calculations are done with antibaryon coupling constants which are given by the G-parity transformation ($g_{\sigma\bar{N}} = g_{\sigma N}$, $g_{\omega\bar{N}} = -g_{\omega N}$) and $SU(3)$ flavor symmetry ($g_{\sigma\bar{\Lambda}} = \frac{2}{3}g_{\sigma\bar{N}}$, $g_{\omega\bar{\Lambda}} = \frac{2}{3}g_{\omega\bar{N}}$). In isosymmetric static systems the scalar and vector potentials for nucleons are expressed as $S = g_{\sigma N}\sigma$ and $V = g_{\omega N}\omega^0$.

3 Nuclear systems containing antibaryons

Now I report on our recent study of antibaryon-doped nuclear systems [4]. Unlike some previous works, we take into account the rearrangement of nuclear structure due to the presence of a real antibaryon. Following the procedure suggested in Ref. [7] and assuming the axial symmetry of the nuclear system, we solve effective Schrödinger equations for nucleons and an antibaryon together with differential equations for mean meson and Coulomb fields. We explicitly take into account the antibaryon contributions to the scalar and vector densities. It is important that antibaryons give a negative contribution to the vector density, while a positive contribution to the scalar density. This leads to increased attraction and decreased repulsion for surrounding nucleons. To maximize attraction, nucleons move to the center of the nucleus, where the antibaryon has its largest occupation probability. This gives rise to a strong local compression of the nucleus and leads to a dramatic rearrangement of its structure.




Figure 1: Single-particle levels in the upper and lower wells for ${}^4\text{He}$ and ${}^{16}\text{O}$ nuclei calculated with the parametrization NL-Z2. Left figures show ordinary nuclei with all levels in the lower well occupied. Right figures show the same nuclei but containing one antiproton (labelled as "+ap") i. e. one hole in the lower well.

These features are illustrated by several examples presented in Figs. 1-3. Figure 1 shows single-particle states in upper and lower wells for ${}^4\text{He}$ and ${}^{16}\text{O}$ nuclei with and without an antiproton. One can clearly see the rearrangement of nuclear structure due to the presence of a hole in the Dirac sea. It is interesting to note that the highest hole level (lowest antiproton level) moves to about zero energy. This means that the whole antiproton rest mass is eaten up by the attractive potential.

Figure 2 shows 3d plots of nucleon density distributions for ${}^{16}\text{O}$ nucleus with and without an antibaryon ($\bar{p}$, $\bar{\Lambda}$) in a lowest energy state. The calculations show that inserting an antiproton into the ${}^{16}\text{O}$ nucleus leads to the increase of central nucleon density by a factor 2-4 depending on the RMF parametrization. Due to a very deep antiproton potential the binding energy of the whole system is increased significantly as compared with 130 MeV for normal ${}^{16}\text{O}$. The calculated binding energies of the $\bar{p}+{}^{16}\text{O}$ system are 830, 1050 and 1160 MeV for the NL-Z2, NL3 and TM1, respectively. Due to this anomalous binding we call such systems super bound nuclei (SBN). In the case of antilambdas we rescale the coupling constants with a factor 2/3 that leads to the binding energy of 560÷700 MeV for the $\bar{\Lambda}+{}^{16}\text{O}$ system.

As second example, we investigate the effect of a single antiproton inserted into the ${}^8\text{Be}$ nucleus. The normal ${}^8\text{Be}$ nucleus is not spherical, exhibiting a clearly visible 2α structure with the ground state deformation $\beta_2 \simeq 1.20$. As seen in Fig. 3, inserting an antiproton in ${}^8\text{Be}$ results in a much less elongated shape ($\beta_2 \simeq 0.23$) and disappearance of its cluster structure. The binding energy increases from 53 MeV to about 700 MeV. Similar, but weaker effects have been predicted [8] for the K^- bound state in the ${}^8\text{Be}$ nucleus.

The calculations have been performed also with reduced antinucleon coupling con-




Figure 2: Sum of neutron and proton densities for ^{16}O (top), ^{16}O with $\bar{\nu}$ (bottom right) and ^{16}O with $\bar{\Lambda}$ (bottom left) calculated with the parametrization NL-Z2.




Figure 3: Nucleon densities for ${}^8\text{Be}$ without (left) and with (right) an antiproton calculated with the parametrization NL3.

stands as compared to the G-parity prescription. We have found that the main conclusions about enhanced binding and considerable compression of $\bar{p}$ -doped nuclei remain valid even when these coupling constants are reduced by factor 3 or so.

4 Antibaryon annihilation in nuclei

The crucial question concerning possible observation of the SBNs is their life time. The main decay channel for such states is the annihilation of antibaryons on surrounding nucleons. The energy available for annihilation of a bound antinucleon equals $Q = 2m_N - B_N - B_{\bar{N}}$, where B_N and $B_{\bar{N}}$ are the corresponding binding energies. In our case this energy is at least by a factor 2 smaller as compared with the vacuum value of $2m_N$. This should lead to a significant suppression of the available phase space and thus to a reduced annihilation rate in medium. We have performed detailed calculations assuming that the annihilation rates into different channels are proportional to the available phase space. All intermediate states with heavy mesons like ρ , ω , η as well as multi-pion channels have been considered. Our conclusion is that decreasing the Q value from 2 GeV to 1 GeV may lead to the reduction of total annihilation rate by factor $20 \div 30$. Then we estimate the SBN life times on the level of 5-25 fm/c which makes their observation feasible. This large margin in the life times is mainly caused by uncertainties in the overlap integral between antinucleon and nucleon scalar densities. Longer life times may be expected for SBNs containing antihyperons. The reason is that, instead of pions, more heavy particles, kaons, are required for their annihilation. We have also analyzed multi-nucleon annihilation channels (Pontecorvo-like reactions) and have found their contribution to be less than 40% of the single-nucleon annihilation.




Figure 4: One-body annihilation leading to sharp lines in the energy spectrum of emitted particles.




Figure 5: Left: compression of a nucleus induced by an antibaryon. Right: collective flow of nuclear fragments resulting from the expansion and disintegration of the nucleus after antibaryon annihilation.

5 Formation and signatures of SBNs

We believe that such exotic antibaryon-nuclear systems can be produced by using antiproton beams of multi-GeV energy, e.g. at the future GSI facility. It is well known that low-energy antiprotons annihilate on the nuclear periphery (at about 5% of the normal density). Since the annihilation cross section drops significantly with energy, a high-energy antiproton can penetrate deeper into the nuclear interior. Then it can be stopped there in an inelastic collision with a nucleon, e. g. via the reaction $A(\bar{p}, N\pi)_{\bar{p}}A'$, leading to the formation of a $\bar{p}$ -doped nucleus. Reactions like $A(\bar{p}, \Lambda)_{\bar{\Lambda}}A'$ can be used to produce a $\bar{\Lambda}$ -doped nuclei. Fast nucleons or lambdas can be used for triggering such events. In order to be captured by a target-nucleus final antibaryons must be slow in the lab frame. Rough estimates of the SBN formation probability in a central $\bar{p}A$ collision give the values $10^{-5} - 10^{-6}$. With the $\bar{p}$ beam luminosity of $2 \cdot 10^{32} \text{ cm}^{-2}\text{s}^{-1}$ planned at GSI this will correspond to the reaction rate from tens to hundreds desired events per second.

Several signatures of SBNs can be used for their experimental observation. First, annihilation of a bound antibaryon can proceed via emission of a single photon, pion or kaon with an energy of about 1 GeV (such annihilation channels are forbidden in vacuum). So one may search for relatively sharp lines, with width of $10 \div 40 \text{ MeV}$, around this energy, emitted isotropically in the SBN rest frame (see Fig. 4). Another signal may come from explosive disintegration of the compressed nucleus after the antibaryon annihilation (see Fig. 5). This process can be observed by measuring radial collective velocities of nuclear fragments.

6 Multi-quark-antiquark clusters

It is interesting to look at the antibaryon-nuclear systems from somewhat different point of view. An antibaryon implanted into a nucleus acts as an attractor for surrounding nucleons. Due to the uncompensated attractive force these nucleons acquire acceleration towards the center. As the result of this inward collective motion the nucleons pile up producing local compression. If this process would be completely elastic it would generate monopole-like oscillations around the compressed SBN state. The maximum compression is reached when the attractive potential energy becomes equal to the compression energy. Simple estimates show that local baryon densities up to 5 times the normal nuclear density may be obtained in this way. It is most likely that the deconfinement transition will occur at this stage and a high-density cloud containing an antibaryon and a few nucleons will appear in the form of a multi-quark-antiquark cluster. One may speculate that the whole ${}^4\text{He}$ or even ${}^{16}\text{O}$ nucleus can be transformed into the quark phase by this mechanism. This process is illustrated in Fig. 6. As shown in ref. [9], an admixture of antiquarks to cold quark matter is energetically favorable. The problem of annihilation is now transferred to the quark level. But the argument concerning the reduction of available phase space due to the entrance-channel nuclear effects should work in this case too. Thus one may hope to produce relatively cold droplets of the quark phase by the inertial compression of nuclear matter initiated by an antibaryon.

I am grateful to T. Bürvenich, W. Greiner and L.M. Satarov for fruitful discussions




Figure 6: Formation of a multi-quark-antiquark cluster at an intermediate stage of the antibaryon annihilation in a ${}^4\text{He}$ nucleus.

and help in the preparation of this talk.

References

- [1] D. Hardtke and STAR Collaboration, Nucl. Phys. **A698** (2002) 671.
- [2] R. Debbe and BRAHMS Collaboration, nucl-ex/0308004.
- [3] N. Auerbach, A.S. Goldhaber, M.B. Johnson, L.D. Miller, A. Picklesimer, Phys. Lett. **B182** (1986) 221.
- [4] T. Bürvenich, I.N. Mishustin, L.M. Satarov, J.A. Maruhn, H. Stöcker, W. Greiner, Phys. Lett. **B542** (2002) 261.
- [5] Y. Sugahara and H. Toki, Nucl. Phys. **A579** (1994) 557.
- [6] M. Bender, K. Rutz, P.-G. Reinhard, J.A. Maruhn, and W. Greiner, Phys. Rev. **C60** (1999) 34304.
- [7] G. Mao, H. Stöcker, and W. Greiner, Int. J. Mod. Phys. **E8** (1999) 389.
- [8] Y. Akaishi and T. Yamazaki, Phys. Rev. **C65** (2002) 044005.
- [9] I.N. Mishustin, L.M. Satarov, H. Stoecker, and W. Greiner, Phys. Rev. **C 59** (1999) 3343.